

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

Lorsban[®]-4E

EPA Reg. No. 62719-220

EPA 24(c) Special Local Need Registration SLN ID-940012

For Distribution and Use Only in the State of Idaho

For Control of Garden Symphylans Infesting Strawberry

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Lorsban[®]-4E insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Lorsban-4E according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Lorsban-4E.

Directions for Use

Preplant Incorporation Treatment

Apply Lorsban-4E in sufficient water to ensure uniform soil coverage and incorporate into the soil in the spring for protection of strawberries during the following year. Use 2 quarts of Lorsban-4E per acre using power-operated ground equipment. On the same day of treatment, incorporate the insecticide into the top 2 to 4 inches of soil using a disc, field cultivator, or equivalent equipment.

Specific Use Precautions:

- Lorsban-4E should not be tank mixed with pesticides, surfactants, or fertilizer formulations unless prior use has shown the combination non-injurious under your current conditions of use. Consult your cooperative agricultural extension service for additional information concerning garden symphylan control practices in your area.

Specific Use Restrictions:

- Apply only as a preplant soil incorporated treatment in the spring for protection of strawberries during the following year.
- Do not make more than 1 application per year of Lorsban-4E.
- Maximum single application rate is 2 lb ai chlorpyrifos per acre.
- **Chemigation:** For uses described in this supplemental labeling, do not apply through any type of irrigation system.
- Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

[®]Trademark of Dow AgroSciences LLC

R062-020

Approved: __/__/__

Replaces D06-062-067

Revisions: Revised paragraph underneath Directions for Use; added "Specific Use Precautions" and revised text underneath; added "Specific Use" before "Restrictions"; added three points.